

## Gel / liquid

**COMPOSITION**

Lidocaine hydrochloride, cetrimide, hypoallergenic/fruit flavors

**PROPERTIES**

**DESENSETIN gel** easily applies and washes out with water. Gel quickly effects on receptors. Thixotropic properties of gel allow to apply it on lower and upper jaws as well as on tongue surface. Spot-on application of gel doesn't spread.

**DESENSETIN liquid** evenly spreads on oral mucosa surface with the help of spraying nozzle, quickly effects on receptors and easily washes out with water. Maximum recommended volume for topical anesthesia is 2 ml. One shot sprinkles 0.12–0.15 ml.

Topical anesthesia starts in 10–15 sec and works for 20–25 minutes, which allows fulfilling of planned treatment procedures.

**INDICATIONS**

Gel and Liquid DESENSETIN are used for topical anesthesia of oral mucosa:

- before injection
- in case of slight injuries
- with gingivectomy
- before scaling
- during teeth cutting
- before extraction of loose teeth
- to reduce gagging with sensitive patients

**CONTRAINDICATIONS**

Drug idiosyncrasy

Use with great care for patients allergic to flavors

Do not use for patients allergic to lidocaine hydrochloride

Do not misuse

**SIDE EFFECTS**

As far as all terms of storing, transporting and application are observed, there are no side effects

**METHOD OF USE**

**Attention!** It is recommended to store material at room temperature (22–24) °C within 1–2 hours before use.

Stir **gel** in jar with dry instrument before use.

Dry oral mucosa with an air jet or cotton ball. Apply **gel** on oral mucosa surface with the help of applicator tip.

Before applying **liquid**, change the transportation cap for dispenser with pump, take off protection cap and put on a spraying nozzle. Apply the liquid with 1–2 press on a bottle pump for anesthesia.

After each application treat the spraying nozzle with disinfectant.

In 10–15 sec planned treatment procedures can be fulfilled. Working time of anesthesia can be prolonged up to 30–40 minutes by reapplication of medication (**gel / liquid**).

Close the materials tightly after use. Being kept open for a long time the anesthetic gel thickens and decreases its properties. At the slight thickening it should be diluted with the distilled water and mixed for the original viscosity.

**PRESENTATION**

Gel (tube)	60 g
Gel (jar)	30 ml (35 g)
Liquid (bottle)	15 ml

**STORAGE**

Store at the temperature from 5 °C to 25 °C  
Keep in dry place  
Tightly close containers immediately after use  
Do not use after expiry date  
Shelf life — 3 years

**MANUFACTURER**

TehnoDent Co. Ltd  
1/6 Beryozovaya str., Severnyi,  
Belgorod region, Russia, 308570  
Tel. +7 (495) 118-28-73  
E-mail: [info@tehnodent.org](mailto:info@tehnodent.org)  
[www.tehnodent.org](http://www.tehnodent.org)

***For professional use only***

1.010.002-01.02\_EN\_002

Rev. 2019-07